

Board Certified Gastroenterologist 1401 S. Beretania St. Suite 320 Honolulu, HI 96814 P (808) 888-0967 F (808) 888-0956

Name :	
value .	

Colonoscopy Preparation

Please follow the instructions precisely to ensure a successful colonoscopy. **Incomplete preparation will lead to a repeat of the procedure and additional out-of-pocket patient cost.** If you need to reschedule, have any questions, or are experiencing difficulties, please call (808)888-0967 Mon-Fri 8:00am-4:00pm. If after office hours or on weekends, please call Physicians Exchange at (808)524-2575 and ask to speak with the Gastroenterologist on call.

Instructional video may be watched on: https://vimeo.com/562638932 or on www.gihealthhawaii.com under the tab "Procedure Preparation"

DAY OF THE	PROCEDURE:			
CHECK-IN TIME:	PROCEDURE TIME:			
LOCATION: LOCATI				
	FIC ENDOSCOPY CENTER 1029 MAKOLU ST, STE H 456-6420			
ONE WEEK BEFORE	YOUR PROCEDURE:			
☐ Please call 808	8-888-0967 to review the instructions and address any questions.			
☆ DO NOT eat popcorn, seeds, corn, or nuts.				
<mark>☆ ***If you tak</mark>	ce blood thinners, please stop oi			
//	*** (days prior to your procedure)			
	**Please do not discontinue your medication until the nurse calls and confirms a			
cl	learance was obtained from your cardiologist.**			

- ☐ Confirm you will have a driver to pick you up after the procedure.
 - The driver must be 18 years or older and present at your discharge.
 - You are REQUIRED to have an adult accompany you to any procedure in which you undergo anesthesia due to the risks and complications that may arise following the procedure.
 - If you do not have a driver, we can arrange to call a Medicab through Charley's Taxi at the patient's expense. Please let the staff know at check-in if this service is needed.

☆ Make sure you have picked up your colonoscopy prep kit.

Please call the office at (808) 888-0967 if we need to resend the prescription.

☐ It is recommended to have petroleum jelly (i.e. Vaseline or Aquaphor) to minimize anal skin irritation.

THREE DAYS BEFORE YOUR PROCEDURE:

- ☆ Eat low-fiber foods.
- ★ Stop taking fiber supplements and anti-diarrheal medications.

LOW FIBER DIET FOR COLONOSCOPY			
✓ Foods that are OK			
White bread, Tofu	Whole wheat bread or pasta		
White rice or noodles	Brown or wild rice		
Plain crackers, vanilla wafers	Whole wheat crackers and rolls		
Skinless cooked potatoes	Raw or undercooked vegetables		
Skinless chicken, turkey, eggs or fish	Tough meat or skinned meat		
Canned fruits (no seeds or skin)	Cereals, granola, cornbread		

ONE DAY BEFORE YOUR PROCEDURE:

- ☆ In the morning, mix your colon preparation with water according to the instructions. Place in the refrigerator to chill. (You must use the solution within 24 hours after mixing.)
- ☆ You may eat breakfast and a light lunch.
- ★ If you take blood pressure medications, aspirin or NSAIDS, you may continue to take them as usual unless otherwise instructed by your physician.
- ★ DIABETIC PATIENTS: No oral medications. INSULIN USERS: take ½ of the usual dose.
- ★ NO SOLID FOODS starting from 1:00PM. until after your procedure
 - **YOU MAY DRINK:**
 - Gatorade, Crystal Light Lemonade, Pedialyte, or Powerade
 - Coffee or Tea (Black Only, Sugar OK)
 - Carbonated or non-carbonated soda
 - Fruit-flavored drinks
 - Apple juice, white cranberry juice or white grape juice
 - Jell-O (gelatin) or popsicles
 - Broth

O DO NOT DRINK THESE LIQUIDS

- Alcohol
- Milk or non-dairy creamer
- Juice with pulp
- Any liquid you cannot see through
- No noodles or vegetables in soup
- Hard Candy

NO RED, ORANGE, OR PURPLE COLORED DRINKS

ONE DAY BEFORE YOUR PROCEDURE:

☆ START DRINKING THE COLON PREPARATION SOLUTION AT 5:00P.M.

- Drink one cup of the solution every 15 minutes until HALF the mixture is gone.
 - *This is the recommended pace. However, if you feel nausea or vomiting, it is perfectly fine to take frequent breaks and drink the solution at your own pace.
- Put the remaining solution in the refrigerator.
- Stay near a toilet as you will have diarrhea, which can start one to five hours after you drink the solution. This can be sudden and last two or more hours after finishing the solution.
- ☆ Drink water often to avoid dehydration. Apply petroleum jelly as needed to anal skin region to alleviate skin irritation.
- ☆ Please do not skip the instructions for the bowel preparation on the day of your colonoscopy due to our body's production of bile overnight.

ΑY	OF THE PROCEDURE:		
HEC	CK-IN TIME:	PROCEDURE TIME:	
	(*NO SMOKING ON THE DAY	OF THE PROCEDURE*)	
☆	On the day of your procedure,	at (5 hours bef	<mark>ore you leave</mark>
	your house) start drinking the	remaining half of the solution.	
☆		vater. Apply petroleum jelly as nee stool is clear or yellow liquid.	eded.
☆	Take your morning medications	at least 2 hours before you leave	for your
	procedure.		
	★ DIABETIC PATIENTS: No.	o oral medications until after the	procedure with
	food. INSULIN USERS: n	o insulin until after the procedur	<mark>e with food</mark>
☆	TWO HOURS BEFORE YOUR (CHECK-IN TIME:BY MOUTH, INCLUDING WATER	_ R, MINT & GUM

= 1 tease teave att jewett y and personal terms at nome.
☐ Please do not wear contact lenses.
☆ Please wear socks to keep your feet warm.
If you have removable dental wear, a denture cup will be provided.
☆ Please bring your I.D. and insurance cards to your appointment.
☆ Please bring a list of all your current medications, including over-the-counter.
☆ Your facility copayment or co-responsibility will be collected on the date of service
unless you made other arrangements. Please bring your preferred form of
payment (credit card, check, or cash). The physician fee and/or pathology fees
will be billed separately and sent to the patient after the insurance has
processed the claim.
Expected duration at the Facility will be between 2-3 hours.
\bigstar Please be aware that there may be unexpected delays from preceding cases.
Though the risk is small, if complications arise, you may be admitted to the
hospital for further observations. Complications may include bleeding, perforation
or adverse reaction to the anesthesia.

COLON CLEANSING TIPS:

1. Drink the solution with a straw to make it easier to tolerate.

D Please leave all jewelry and personal items at home

- 2. If you experience nausea or vomiting, give yourself a 30 minute break and then try to drink the prep solution again.
- 3. You may experience cramps until all the stool has flushed from your colon, which may take 2-4 hours or longer.
- 4. Anal skin irritation or flare of hemorrhoid inflammation may occur. You may use over-the-counter remedies, such as hydrocortisone cream, baby wipes, Vaseline, or TUCKS pads to improve comfort. Avoid products containing alcohol.

☆ If you develop symptoms such as cold, fever, persistent coughs, or runny nose, please call (808) 888-0967 to reschedule your appointment. The facility will contact you 1-2 days prior to confirm your appointment and to complete a COVD-19 questionnaire.

MEDICATION LIST:

(*PLEASE COMPLETE THE FORM AND BRING TO THE FACILITY CENTER ON THE DAY OF YOUR PROCEDURE*)

SOURCE OF INFORMATION: PATIENT FAMILY OTHE
--

ALLERGIES	REACTION	ALLERGIES	REACTION

MEDICATION NAME	DOSE	FREQUENCY	ROUTE	REASON	LAST TAKEN
1.(example) Aspirin	81 mg	once a day	oral	Blood thinner	5am today
2.					
3.					
4.					
5.					
6.					
7.					
8.					
9.					
10.					
11.					
12.					
13.					
14.					