



Board Certified Gastroenterologist
1401 S. Beretania St. Suite 320
Honolulu, HI 96814
P (808) 888-0967 F (808) 888-0956

Name:

SUPREP Colonoscopy Preparation

Please follow the instructions precisely to ensure a successful colonoscopy. **Incomplete preparation will lead to a repeat of the procedure and additional out-of-pocket patient cost.** If you need to reschedule, have any questions, or are experiencing difficulties, please call **(808)888-0967** Mon-Fri 8:00am-4:00pm. If after office hours or on weekends, please call Physicians Exchange at (808)524-2575 and ask to speak with the Gastroenterologist on call.

DAY OF THE PROCEDURE:

CHECK-IN TIME:

PROCEDURE TIME:

LOCATION: ENDOSCOPY INSTITUTE OF HI 1401 S. BERETANIA ST,
Suite 200, 2nd Floor (808)312-6700

PACIFIC ENDOSCOPY CENTER 1029 MAKOLU ST, **STE H**
(808)456-6420

ONE WEEK BEFORE YOUR PROCEDURE:

- Please call 808-888-0967 to review the instructions and address any questions.
- DO NOT** eat popcorn, seeds, corn, or nuts.
- If you take blood thinners, please STOP _____ on _____**

****Please do not discontinue your medication until the nurse calls and confirms a clearance was obtained from your cardiologist.****

- ❑ Confirm you will have a driver to pick you up after the procedure.
 - The driver must be 18 years or older and present at your discharge.
 - You are REQUIRED to have an adult accompany you to any procedure in which you undergo anesthesia due to the risks and complications that may arise following the procedure.
 - If you do not have a driver, we can arrange to call a Medicab through Charley's Taxi at the patient's expense. Please let the staff know at check-in if this service is needed.

★ Make sure you have your colonoscopy preparation kit. Please call the office at (808)888-0967 if we need to resend the prescription.

★ Please also purchase petroleum jelly (i.e. Vaseline or Aquaphor) to alleviate anal skin irritation.

THREE DAYS BEFORE YOUR PROCEDURE:

★ Eat low-fiber foods.

★ Stop taking fiber supplements and anti-diarrheal medications.

LOW FIBER DIET FOR COLONOSCOPY	
✓ Foods that are OK	⊘ Foods to AVOID
White bread, Tofu	Whole wheat bread or pasta
White rice or noodles	Brown or wild rice
Plain crackers, vanilla wafers	Whole wheat crackers and rolls
Skinless cooked potatoes	Raw or undercooked vegetables
Skinless chicken, turkey, eggs or fish	Tough meat or skinned meat
Canned fruits (no seeds or skin)	Cereals, granola, cornbread

ONE DAY BEFORE YOUR PROCEDURE:

☆ You will be on **CLEAR LIQUID DIET ALL DAY** starting when you wake up.

☆ If you take blood pressure medications, aspirin or NSAIDS, you may continue to take them as usual unless otherwise instructed by your physician.

☆ **DIABETIC PATIENTS: No oral medications. INSULIN USERS: take ½ of the usual dose**

<p>✓ YOU MAY DRINK THESE CLEAR LIQUIDS</p> <ul style="list-style-type: none">● Gatorade, Crystal Light Lemonade, Pedialyte, or Powerade● Coffee or Tea (Black Only, Sugar OK)● Carbonated or non-carbonated soda● Fruit-flavored drinks● Apple juice, white cranberry juice or white grape juice● Jell-O (gelatin) or popsicles● Broth	<p>⊘ DO NOT DRINK THESE LIQUIDS</p> <ul style="list-style-type: none">● Alcohol● Milk or non-dairy creamer● Juice with pulp● Any liquid you cannot see through● No noodles or vegetables in soup● Hard Candy <p>***NO RED, ORANGE, OR PURPLE COLORED DRINKS***</p>
--	--

ONE DAY BEFORE YOUR PROCEDURE:

☆ **START DRINKING THE COLON PREPARATION SOLUTION AT 5:00P.M.**

- Pour one (1) 6-ounce bottle of SUPREP liquid into the mixing container. Then add drinking water to the 16-ounce line on the container and mix. Drink ALL the liquid. Then drink two (2) MORE 16-ounce containers of **WATER** over the next hour. ****This is the recommended pace. However, if you feel nausea or vomiting, it is perfectly fine to take frequent breaks and drink the solution at your own pace.***
- Stay near a toilet as you will have diarrhea, which can start one to five hours after you start the solution. This can be sudden and may last two or more hours after finishing the solution.
- Apply petroleum jelly as needed to alleviate anal skin irritation.
- ☆ Drink water often to avoid dehydration.
- ☆ Please do not skip the instructions for the bowel preparation on the day of your colonoscopy due to our body's production of bile overnight.

DAY OF THE PROCEDURE:

CHECK-IN TIME:

PROCEDURE TIME:

(*NO SMOKING ON THE DAY OF THE PROCEDURE*)

☆ On the day of your procedure, at _____ (5 HOURS before you leave home for the procedure) Pour one (1) 6-ounce bottle of SUPREP liquid into the mixing container. Then add drinking water to the 16-ounce line on the container and mix. Drink ALL the liquid. Then drink two (2) MORE 16-ounce containers of **WATER** over the next hour.

★ You are ready when your stool is clear or yellow liquid

★ Apply petroleum jelly as needed to alleviate anal skin irritation.

☆ Take your morning medications at least 2 hours before you leave for your procedure.

● **DIABETIC PATIENTS: No oral medications until after the procedure with food. INSULIN USERS: no insulin until after the procedure with food.**

☆ TWO HOURS BEFORE YOUR CHECK-IN TIME: _____
PLEASE STOP **ANY** INTAKE BY MOUTH, INCLUDING WATER, MINT & GUM

Please leave all jewelry and personal items at home.

Please do not wear contact lenses.

☆ Please wear socks to keep your feet warm.

If you have removable dental wear, a denture cup will be provided.

☆ Please bring your I.D. and insurance cards to your appointment.

☆ Please bring a list of all your current medications, including over-the-counter.

☆ Your facility copayment or co-responsibility will be collected on the date of service unless you made other arrangements. Please bring your preferred form of payment (credit card, check, or cash). **The physician fee and/or pathology fees will be billed separately and sent to the patient after the insurance has processed the claim.**

Expected duration at the Facility will be between 2-3 hours.

★ Please be aware that there may be unexpected delays from preceding cases.

- ❑ Though the risk is small, if complications arise, you may be admitted to the hospital for further observations. Complications may include bleeding, perforation, or adverse reaction to the anesthesia

COLON CLEANSING TIPS:

1. Drink the solution with a straw to make it easier to tolerate.
2. If you experience nausea or vomiting, give yourself a 30 minute break and then try to drink the prep solution again.
3. You may experience cramps until all the stool has flushed from your colon, which may take 2-4 hours or longer.
4. Anal skin irritation or flare of hemorrhoid inflammation may occur. You may use over-the-counter remedies, such as hydrocortisone cream, baby wipes, Vaseline, or TUCKS pads to improve comfort. Avoid products containing alcohol.

★ If you develop symptoms such as cold, fever, persistent coughs, or runny nose, please call **(808) 888-0967** to reschedule your appointment. The facility will contact you 1-2 days prior to confirm your appointment and to complete a COVID-19 questionnaire.

MEDICATION LIST:

(*PLEASE COMPLETE THE FORM AND BRING TO THE FACILITY CENTER ON THE DAY OF YOUR PROCEDURE*)

SOURCE OF INFORMATION: _____ PATIENT _____ FAMILY _____ OTHER

ALLERGIES	REACTION	ALLERGIES	REACTION

MEDICATION NAME	DOSE	FREQUENCY	ROUTE	REASON	LAST TAKEN
1.					
2.					
3.					
4.					
5.					
6.					
7.					
8.					
9.					
10.					
11.					
12.					